

Contribution to differential π^0 and γ_{dir} modification in small systems from color fluctuation effects

Dennis V. Perepelitsa*
University of Colorado Boulder
Boulder, CO 80309

A major complication in the search for jet quenching in proton– or deuteron–nucleus collision systems is the presence of physical effects which influence the experimental determination of collision centrality in the presence of a hard process. For example, in the proton color fluctuation picture, protons with a large Bjorken- x ($x \gtrsim 0.1$) parton interact more weakly with the nucleons in the nucleus, leading to a smaller (larger) than expected yield in large (small) activity events.

A recent measurement by PHENIX compared the yield of neutral pion and direct photon production in $d+\text{Au}$ collisions, under the argument that the photon yields correct for such biases, and the difference between the two species is thus attributable to final-state effects (i.e., jet quenching). The main finding suggests a significant degree of jet quenching for hard processes in small systems.

In this paper, I argue that the particular photon and pion events selected by PHENIX arise from proton configurations with significantly different Bjorken- x distributions, and thus are subject to different magnitudes of modification in the color fluctuation model. Using the results of a previous global analysis of RHIC and LHC data, I show that potentially all of the pion-to-photon difference in PHENIX data can be described by a proton color fluctuation picture at a quantitative level before any additional physics from final-state effects is required. This finding reconciles the interpretation of the PHENIX measurement with others at RHIC and LHC, which have found no observable evidence for jet quenching in small systems.

I. INTRODUCTION

Measurements of hard processes in relativistic proton- or deuteron-nucleus ($p/d+\text{A}$) at the Relativistic Heavy Ion Collider (RHIC) and the Large Hadron Collider (LHC) have a number of scientific purposes [1, 2], including a precision determination of the parton densities in nuclei (see [3] for a recent example), constraints on dynamical processes in the initial state of the cold nucleus (such as those observed at lower energies [4]), and studies of the general interplay between soft and hard processes in collisions involving nuclei. Another key interest, given the robust experimental signatures of Quark Gluon Plasma-like behavior in these “small” systems [5–9], is the search for evidence of final-state interactions between the hard-scattered parton and the dilute system. If the parton shower is modified in the final state as it propagates through the produced system, it may lead to a number of effects, the most direct of which is the decreased production of jets or hadrons at fixed transverse momentum (p_T), i.e., jet quenching [10–12].

By analogy to large, nucleus-nucleus collision systems, one wants to search in the most “central”, or highest-activity, collisions where the produced system is largest and longest-lived. Unfortunately, the experimental selection on centrality in small systems is challenging due to strong auto-correlation biases and non-trivial physics effects, with magnitudes larger than that required for a precision constraint on jet quenching effects. For events with a generic hard process, the selection is sensitive

to upward multiplicity fluctuations (multiplicity vetoes), leading to an apparently enhanced yield in central (decreased yield in peripheral) collisions compared to that expected from the estimated number of nucleon–nucleon collisions, $\langle N_{\text{coll}} \rangle$, in the events [13–16].

As a separate phenomenon, in extreme kinematic regions characterized by Bjorken- $x \gtrsim 0.1$, the pattern reverses, with a significant depletion (enhancement) in the high-activity central (low-activity peripheral) events, as directly observed in measurements by ATLAS [17], CMS [18], STAR [19], and PHENIX [20] in $p/d+\text{A}$ events. In a recent measurement by ATLAS [21], the produced dijet pair was used to estimate the parton-level kinematics in each event, strongly suggesting that the modifications follow a universal pattern in the Bjorken- x of the proton, with their magnitude systematically increasing with x . A particular physics interpretation of these observations in data is the QCD color fluctuation model [22, 23], which proposes that proton configurations with a large- x parton interact more weakly with nucleons in the nucleus than configuration-averaged protons, thus leading to a decreased centrality signal and the particular pattern of x -dependent modifications described above.

To get around these challenges, experimental searches of jet quenching in $p/d+\text{A}$ collisions have therefore been performed in centrality-averaged (minimum-bias) collisions [24–26], or using techniques such as jet–hadron correlations [27, 28] and di-jet asymmetries [18, 19] in central events, all giving null results for a detectable jet quenching-like signature. A recent PHENIX measurement [29] in 200 GeV $d+\text{Au}$ collisions used a different technique and reported the relative yields of neutral pions (π^0 's) to those of direct photons (γ_{dir} 's), under the argument that both are subject to the same cen-

* dvp@colorado.edu

trality bias effects, and thus any difference in the degree of modification should be attributed to centrality-dependent final-state effects (i.e., which would not be felt by the photons). The PHENIX measurement reports the quantity $R_{\text{dAu,EXP}}$, defined as the double ratio of the nuclear modification factors for π^0 's and γ_{dir} , $R_{\text{dAu},\pi^0}/R_{\text{dAu},\gamma_{\text{dir}}}$, i.e., conceptually defining $R_{\text{dAu},\gamma_{\text{dir}}}$ to be unity for all centrality selections. The measurement reports $R_{\text{dAu,EXP}}$ in the most central 0–5% of $d+\text{Au}$ events as $0.77 \pm 0.03(\text{stat}) \pm 0.13(\text{syst})$, where the latter is dominated by a global uncertainty due to the normalization of the γ_{dir} measurement. If interpreted as a jet quenching effect, the data are challenging to understand given the extensive null results described above. For example, the PHENIX data would require a π^0 p_{T} spectrum shift of $\delta p_{\text{T}} \approx -0.21$ GeV in the most central events at RHIC [30], corresponding to a fractional energy loss of $\delta p_{\text{T}}/p_{\text{T}} \approx 1\text{--}2\%$ in $d+\text{Au}$ events, whereas the measurement by ATLAS in Ref. [28] sets an exclusion limit at 90% confidence level of $\delta p_{\text{T}}/p_{\text{T}} < 1.4\%$ for charged hadrons in $p+\text{Pb}$ collisions at the LHC. However, these two measurements probe partons with different initial energies and collisions with different energies and geometries, and thus may not be in explicit tension.

Notably, the π^0 and γ_{dir} were measured in the same p_{T} range of 7.5–18 GeV which, since the π^0 carries only a fraction of the tree-level parton p_{T} , actually corresponds to different Bjorken- x ranges. Thus, the measurement technique defined by PHENIX is directly sensitive to the x -dependent color fluctuation effects described above, in addition to any jet quenching effects. In particular, the strength of the effect changes quickly with x in the region of the measurement, and is thus only partially “calibrated out” with the γ_{dir} baseline. In this paper, I argue that the color fluctuation effect, as calculated using the results of Ref. [23] with no additional model modifications or re-tuning to new data, can potentially explain the full centrality dependence of the observable defined by PHENIX. When accounting for it, the possible magnitude of any jet quenching effect is significantly smaller, reconciling the interpretation of the PHENIX measurement with the established constraints on jet quenching by the other RHIC and LHC experiments.

II. METHOD

The PYTHIA Monte Carlo event generator [31] was used to simulate the nucleon–nucleon sub-collisions in $d+\text{Au}$ collisions which produce π^0 's and γ_{dir} 's in the PHENIX kinematic selections [29], and determine the distribution of Bjorken- x values for these events. The specific selections are $p_{\text{T}} = 7.5\text{--}18$ GeV and $|\eta| < 0.35$ for both species. PYTHIA version 8.307 was used to generate events at $\sqrt{s} = 200$ GeV, with a mixture of proton–proton, proton–neutron, and neutron–neutron collisions appropriate for $d+\text{Au}$. PYTHIA was configured with all HardQCD processes and a $\hat{p}_{\text{T,min}}$ (minimum parton–




FIG. 1. Distribution in PYTHIA8 of the values of Bjorken- x in the deuteron in 200 GeV $d+\text{Au}$ collisions, for events producing π^0 's (black) or γ_{dir} 's (red) in the kinematic selection used by PHENIX in Ref. [29].

parton transverse momentum exchange) of 5 GeV for the π^0 case. For the direct photon case, PYTHIA was configured with all PromptPhoton processes and photons which are the result of hadron decays were rejected (i.e., only those radiated by a quark were selected). This $\hat{p}_{\text{T,min}}$ threshold was checked that it does not introduce a kinematic bias in the final-state p_{T} range of interest. One of the beam hadrons was chosen to represent the proton or neutron in the deuteron of a $d+\text{Au}$ collision and its Bjorken- x value, defined in PYTHIA as the x at which the PDF in the beam is defined [31], was recorded.

Figure 1 shows the distribution of Bjorken- x values in the deuteron for events with a π^0 or γ_{dir} within the PHENIX kinematic selection. For γ_{dir} , the distribution is peaked near $x \sim 7.5/100 = 0.075$, where 7.5 GeV and 100 GeV are the minimum γ_{dir} p_{T} threshold and the beam energy, respectively, with a small tail of contributions from events with higher x . For π^0 , the distribution has a significantly harder tail to large- x values, reflecting the fact that they are produced by the fragmentation of hard scattered partons and thus carry only a fraction of their energy. Notably, the average x value is more than 50% larger for π^0 - than γ_{dir} -producing events, with 28% (9%) of the π^0 's (γ_{dir} 's) produced in events with $x > 0.2$. In principle, the magnitude of the color fluctuation effect is expected to be different depending on whether the high- x parton in the deuteron is a quark or a gluon. However, in these kinematics, the quark/gluon fraction is similar in the π^0 and γ_{dir} events and thus the difference in their x distributions is the main distinction.

The distributions of x values in Fig. 1 were then folded with the results of the color fluctuation model in Ref. [23] to produce a prediction for the results of the PHENIX measurement. First, the color fluctuation model paper primarily reports R_{CP} , the ratio of nuclear modification in central to peripheral events, which must be converted to the $R_{\text{dAu,EXP}}$ and other observables reported in the




FIG. 2. Calculated nuclear modification factor R_{dAu} within the color fluctuation model as a function of the average number of colliding nucleon pairs, $\langle N_{coll} \rangle$. Calculations are shown for π^0 's (black) and γ_{dir} 's (red) matching the PHENIX kinematic selection. Since this model does not include any overall modification, the minimum-bias R_{dAu} is unity by construction.

recent PHENIX paper. This is accomplished using the detailed information provided in an earlier PHENIX publication [13] on the geometric properties (e.g. the average number of colliding nucleon–nucleon pairs $\langle N_{coll} \rangle$) of centrality-selected $d+Au$ events. Importantly, while the results here are presented as a function of the average $\langle N_{coll} \rangle$, the calculations in Ref. [23] on which they are based were determined using the distribution of the N_{coll} values in each centrality bin. Second, the color fluctuation model presents results in exclusive Bjorken- x bins, which are then combined in a weighted average with the x distributions for π^0 and γ_{dir} events above. I highlight that the parameters of this model, determined using previous RHIC and LHC data, have not been adjusted or updated before the comparison to the PHENIX data of interest.

Notably, the color fluctuation model in Ref. [23] does not itself include any centrality-independent modification, i.e., R_{dAu} for minimum-bias events is unity by construction. However, there are expected to be small modifications of the minimum-bias rate for both species due to the nuclear parton densities [3] and the “isospin effect” for direct photon production from the valence quark region [32, 33]. Given the specific interest in the centrality dependence of the observable and the small magnitude of the minimum-bias-averaged effects compared to the large overall normalization uncertainties in PHENIX data, I do not model these effects explicitly in this work. If such effects were included, they would result in a global rescaling of all calculated R_{dAu} points in the same direction.

Fig. 2 shows the calculated nuclear modification factors, R_{dAu} , within the color fluctuation model, applied to the π^0 and γ_{dir} events in the PHENIX kinematics selection, and within the PHENIX centrality intervals.




FIG. 3. Comparison of the nuclear modification factor R_{dAu} for direct photons as a function of $\langle N_{coll} \rangle$, showing the measurement in PHENIX data (black points) and the calculation from the color fluctuation model (red horizontal lines). The vertical bars and boxes around the data points indicate the statistical uncertainties and the overall 16.5% normalization uncertainty, respectively.

Since the minimum-bias R_{dAu} is unity by construction (see above), the modifications appear as a relative enhancement/suppression pattern around unity in peripheral/central events. Importantly, the magnitude of the relative modifications is stronger for π^0 's than for γ_{dir} 's, reflecting the different x distributions in Fig. 1. Indeed, in Ref. [23], the quantity $\langle \lambda(x) \rangle$, which describes the x -dependent weakening of the inelastic nucleon–nucleon interaction strength, falls rapidly from ≈ 0.8 to ≈ 0.6 in this Bjorken- x region, resulting in the difference seen here.

III. COMPARISON TO DATA

Fig. 3 compares the calculated R_{dAu} for γ_{dir} integrated over the $p_T = 7.5\text{--}18$ GeV range within the color fluctuation model to that measured by PHENIX, where the quantity is styled as $(Y_{dAu}^{\gamma_{dir}}/Y_{pp}^{\gamma_{dir}})/N_{coll}^{Glauber}$. For this comparison, the measured minimum-bias R_{dAu} for γ_{dir} is essentially unity, and so no overall rescaling of the model calculation is applied. The model gives a good description of the modest centrality dependence of the data, with a relative enhancement and suppression pattern in going from peripheral to central events. As a reminder, the γ_{dir} selection typically results in events with $x \approx 0.11$, and thus the impact of the color fluctuation physics on this observable is modest in the x range relevant for γ_{dir} production.

Fig. 4 compares the measured and calculated “double ratio” of the R_{dAu} for π^0 's to that for γ_{dir} 's, as a function of $\langle N_{coll} \rangle$, which is the main result of the PHENIX analysis. The dashed red line shows the result of the color fluctuation model, under the assumption that the




FIG. 4. Comparison of the ratio of nuclear modification factors between that for neutral pions and direct photons, $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$, as a function of $\langle N_{coll} \rangle$, showing the measurement in PHENIX data (black points) and the calculation from the color fluctuation model (red lines). The vertical bars and boxes around the data points indicate the statistical uncertainties and the overall 16.5% normalization uncertainty, respectively. The dashed red line assumes that R_{dAu} in minimum-bias events for both π^0 's and γ_{dir} 's is unity. The solid red line adjusts the calculation by a factor of 0.9, to better match the actually measured $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$ in minimum-bias events by PHENIX.

R_{dAu} in minimum-bias events is unity for both species. The calculation gives a good, quantitative description of the data within its significant normalization uncertainty, including the slow decrease of the double ratio with increasing $\langle N_{coll} \rangle$.

In the PHENIX analysis, the minimum bias $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$ is measured to be $0.92 \pm 0.02(\text{stat}) \pm 0.15(\text{syst})$. Ref. [29] notes that this value is consistent with unity within the large overall scale uncertainty, which is dominated by uncertainties in the π^0 and γ_{dir} $p+p$ references. Thus, the data-model comparison here has a significant degree of freedom in the overall normalization of the data. The best match between the model and the central values of the data is achieved by adjusting the model down by a factor 0.90, shown as a solid red line in Fig. 4. I note that this factor is close to the measured minimum-bias $R_{dAu}(\pi^0)/R_{dAu}(\gamma_{dir})$ value of 0.92, and is well within the normalization uncertainty of the data. This may alternatively be thought of as scaling the data up by approximately two thirds of its stated global uncertainty. With this normalization, the model now provides an excellent description of the PHENIX data in detail. While not explored here, I note that taking into account the uncertainties in the model parameters themselves [23] may further improve the data-model agreement. Thus, no other physics effects, such as a significant centrality-dependent jet quenching, are needed to describe the PHENIX data.

In the future, several experimental avenues may help to separate the contribution of the color fluctuation effect from searches for the jet quenching physics of interest. First, one may design the kinematic selections with an aim to match the Bjorken- x ranges accessed by the hadron/jet and photon processes, and thus cancel the impact of the color fluctuation effect. Second, as motivated in Ref. [34], the two effects are expected to have different sensitivities to changing the projectile, such as in $p+Au$ or $^3\text{He}+Au$ data previously recorded at RHIC. This has been performed for π^0 's in Ref. [35], but not yet for γ_{dir} . Third, one could attempt a centrality selection based on measuring spectator neutrons in a zero-degree calorimeter (e.g. as used in Refs [27, 28]), which may better isolate the underlying geometry and not be sensitive to physics effects which manifest as a bias on the produced multiplicity. Fourth, oxygen-oxygen (O+O) collisions at RHIC and the LHC [36] would allow for the study of jet quenching in a small system, in which a single weakly-interacting nucleon in the ^{16}O nucleus will have a much smaller impact on the overall centrality signal.

IV. CONCLUSION

This paper examines a recent PHENIX measurement which found a different degree of centrality-dependent modification for π^0 's and γ_{dir} 's in $d+Au$ collisions at RHIC. The measurement strategy is ostensibly chosen to calibrate out any centrality-dependent biases with the γ_{dir} measurement. However, I show that the particular π^0 and γ_{dir} kinematic selections used in the measurement select events which arise from different distributions of Bjorken- x values. As such, the PHENIX measurement is directly sensitive to the physics effects described by the color fluctuation model, in which centrality-dependent modifications (which do not arise from jet quenching) systematically increase with x . Using the model parameters determined from RHIC and LHC measurements reported ten years ago, without any post-diction updating, I show that a straightforward application of the model to the PHENIX kinematics gives a good description of the experimental data. After accounting for the possible role of color fluctuation effects, the evidence for any remaining final-state effects, such as from jet quenching, is significantly more limited. This finding reconciles the interpretation of the PHENIX data with that from other measurements at RHIC and LHC, which have set stringent limits on the possible amount of parton energy loss in small collision systems.

ACKNOWLEDGMENTS

DVP acknowledges John Lajoie, Riccardo Longo, Jamie Nagle, Peter Steinberg, and Mark Strikman for useful discussions and input on an early draft of this paper. DVP's work is supported by the U.S. Department of Energy, grant DE-FG02-03ER41244.

-
- [1] C. A. Salgado *et al.*, Proton-Nucleus Collisions at the LHC: Scientific Opportunities and Requirements, *J. Phys. G* **39**, 015010 (2012), arXiv:1105.3919 [hep-ph].
- [2] C. A. Salgado and J. P. Wessels, Proton-Lead Collisions at the CERN LHC, *Ann. Rev. Nucl. Part. Sci.* **66**, 449 (2016).
- [3] K. J. Eskola, P. Paakkinen, H. Paukkunen, and C. A. Salgado, EPPS21: a global QCD analysis of nuclear PDFs, *Eur. Phys. J. C* **82**, 413 (2022), arXiv:2112.12462 [hep-ph].
- [4] S. J. Paul *et al.* (CLAS), Observation of Azimuth-Dependent Suppression of Hadron Pairs in Electron Scattering off Nuclei, *Phys. Rev. Lett.* **129**, 182501 (2022), arXiv:2207.06682 [nucl-ex].
- [5] J. L. Nagle and W. A. Zajc, Small System Collectivity in Relativistic Hadronic and Nuclear Collisions, *Ann. Rev. Nucl. Part. Sci.* **68**, 211 (2018), arXiv:1801.03477 [nucl-ex].
- [6] R. D. Weller and P. Romatschke, One fluid to rule them all: viscous hydrodynamic description of event-by-event central p+p, p+Pb and Pb+Pb collisions at $\sqrt{s} = 5.02$ TeV, *Phys. Lett. B* **774**, 351 (2017), arXiv:1701.07145 [nucl-th].
- [7] J. Adam *et al.* (ALICE), Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions, *Nature Phys.* **13**, 535 (2017), arXiv:1606.07424 [nucl-ex].
- [8] A. M. Sirunyan *et al.* (CMS), Elliptic flow of charm and strange hadrons in high-multiplicity pPb collisions at $\sqrt{s_{NN}} = 8.16$ TeV, *Phys. Rev. Lett.* **121**, 082301 (2018), arXiv:1804.09767 [hep-ex].
- [9] G. Aad *et al.* (ATLAS), Transverse momentum and process dependent azimuthal anisotropies in $\sqrt{s_{NN}} = 8.16$ TeV p+Pb collisions with the ATLAS detector, *Eur. Phys. J. C* **80**, 73 (2020), arXiv:1910.13978 [nucl-ex].
- [10] K. Tywoniuk, Is there jet quenching in pPb?, *Nucl. Phys. A* **926**, 85 (2014).
- [11] A. Huss, A. Kurkela, A. Mazeliauskas, R. Paatelainen, W. van der Schee, and U. A. Wiedemann, Discovering Partonic Rescattering in Light Nucleus Collisions, *Phys. Rev. Lett.* **126**, 192301 (2021), arXiv:2007.13754 [hep-ph].
- [12] D. V. Perepelitsa, Connection between soft and hard probes of small collision systems at RHIC and LHC, *EPJ Web Conf.* **235**, 04003 (2020), arXiv:2005.05981 [nucl-ex].
- [13] A. Adare *et al.* (PHENIX), Centrality categorization for $R_{p(d)+A}$ in high-energy collisions, *Phys. Rev. C* **90**, 034902 (2014), arXiv:1310.4793 [nucl-ex].
- [14] J. Adam *et al.* (ALICE), Centrality dependence of particle production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. C* **91**, 064905 (2015), arXiv:1412.6828 [nucl-ex].
- [15] D. V. Perepelitsa and P. A. Steinberg, Calculation of centrality bias factors in p+A collisions based on a positive correlation of hard process yields with underlying event activity, (2014), arXiv:1412.0976 [nucl-ex].
- [16] C. Loizides and A. Morsch, Absence of jet quenching in peripheral nucleus-nucleus collisions, *Phys. Lett. B* **773**, 408 (2017), arXiv:1705.08856 [nucl-ex].
- [17] G. Aad *et al.* (ATLAS), Centrality and rapidity dependence of inclusive jet production in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector, *Phys. Lett. B* **748**, 392 (2015), arXiv:1412.4092 [hep-ex].
- [18] S. Chatrchyan *et al.* (CMS), Studies of dijet transverse momentum balance and pseudorapidity distributions in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Eur. Phys. J. C* **74**, 2951 (2014), arXiv:1401.4433 [nucl-ex].
- [19] Event activity correlations and jet measurements in p+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, (2024), arXiv:2404.08784 [nucl-ex].
- [20] A. Adare *et al.* (PHENIX), Centrality-dependent modification of jet-production rates in deuteron-gold collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* **116**, 122301 (2016), arXiv:1509.04657 [nucl-ex].
- [21] G. Aad *et al.* (ATLAS), Measurement of the Centrality Dependence of the Dijet Yield in p+Pb Collisions at $s_{NN} = 8.16$ TeV with the ATLAS Detector, *Phys. Rev. Lett.* **132**, 102301 (2024), arXiv:2309.00033 [nucl-ex].
- [22] M. Alvioli, B. A. Cole, L. Frankfurt, D. V. Perepelitsa, and M. Strikman, Evidence for x -dependent proton color fluctuations in pA collisions at the CERN Large Hadron Collider, *Phys. Rev. C* **93**, 011902 (2016), arXiv:1409.7381 [hep-ph].
- [23] M. Alvioli, L. Frankfurt, D. Perepelitsa, and M. Strikman, Global analysis of color fluctuation effects in proton- and deuteron-nucleus collisions at RHIC and the LHC, *Phys. Rev. D* **98**, 071502 (2018), arXiv:1709.04993 [hep-ph].
- [24] V. Khachatryan *et al.* (CMS), Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *JHEP* **04**, 039, arXiv:1611.01664 [nucl-ex].
- [25] S. Acharya *et al.* (ALICE), Nuclear modification factor of light neutral-meson spectra up to high transverse momentum in p-Pb collisions at $s_{NN} = 8.16$ TeV, *Phys. Lett. B* **827**, 136943 (2022), arXiv:2104.03116 [nucl-ex].
- [26] G. Aad *et al.* (ATLAS), Charged-hadron production in pp, p+Pb, Pb+Pb, and Xe+Xe collisions at $\sqrt{s_{NN}} = 5$ TeV with the ATLAS detector at the LHC, *JHEP* **07**, 074, arXiv:2211.15257 [hep-ex].
- [27] S. Acharya *et al.* (ALICE), Constraints on jet quenching in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured by the event-activity dependence of semi-inclusive hadron-jet distributions, *Phys. Lett. B* **783**, 95 (2018), arXiv:1712.05603 [nucl-ex].
- [28] G. Aad *et al.* (ATLAS), Strong Constraints on Jet Quenching in Centrality-Dependent p+Pb Collisions at 5.02 TeV from ATLAS, *Phys. Rev. Lett.* **131**, 072301 (2023), arXiv:2206.01138 [nucl-ex].
- [29] N. J. Abdulameer *et al.* (PHENIX), Disentangling centrality bias and final-state effects in the production of high- $p_T \pi^0$ using direct γ in d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, (2023), arXiv:2303.12899 [nucl-ex].
- [30] A. Drees, PHENIX Overview (2023), 7th International Conference on the Initial Stages in High-Energy Nuclear Collisions.
- [31] C. Bierlich *et al.*, A comprehensive guide to the physics and usage of PYTHIA 8.3, *SciPost Phys. Codeb.* **2022**, 8 (2022), arXiv:2203.11601 [hep-ph].

- [32] A. Adare *et al.* (PHENIX), Direct photon production in $d+Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C **87**, 054907 (2013), arXiv:1208.1234 [nucl-ex].
- [33] M. Aaboud *et al.* (ATLAS), Measurement of prompt photon production in $\sqrt{s_{NN}} = 8.16$ TeV $p+Pb$ collisions with ATLAS, Phys. Lett. B **796**, 230 (2019), arXiv:1903.02209 [nucl-ex].
- [34] D. McGlinchey, J. L. Nagle, and D. V. Perepelitsa, Consequences of high- x proton size fluctuations in small collision systems at $\sqrt{s_{NN}} = 200$ GeV, Phys. Rev. C **94**, 024915 (2016), arXiv:1603.06607 [nucl-th].
- [35] U. A. Acharya *et al.* (PHENIX), Systematic study of nuclear effects in $p+Al$, $p+Au$, $d+Au$, and ^3He+Au collisions at $\sqrt{s_{NN}} = 200$ GeV using π^0 production, Phys. Rev. C **105**, 064902 (2022), arXiv:2111.05756 [nucl-ex].
- [36] J. Brewer, A. Mazeliauskas, and W. van der Schee, Opportunities of OO and pO collisions at the LHC, in *Opportunities of OO and pO collisions at the LHC* (2021) arXiv:2103.01939 [hep-ph].